

Fill out this medical information card with your doctor's contact information. You can keep it with you while on BLINCYTO® treatment.



Fold

<p>About me:</p> <p>_____</p> <p>Name</p> <p>_____</p> <p>Phone</p> <p>My parent or caregiver:</p> <p>_____</p> <p>Name</p> <p>_____</p> <p>Phone</p>	<p>Learn more at www.BLINCYTO.com</p> <p>_____</p> <p>Phone</p> <p>_____</p> <p>Name</p> <p>_____</p> <p>Name</p> <p>About my doctor:</p>
<p>AMGEN®</p> <p>BLINCYTO® is a registered trademark of Amgen Inc. © 2018 Amgen Inc. All rights reserved. USA-103-80105 10/18</p> <p>Fill out this medical information card. You can keep your important health care contact information in one place here. There is also information on some possible side effects.</p>	<p>Cytokine release syndrome and infusion reactions</p> <p>Symptoms may include:</p> <ul style="list-style-type: none"> • Fever • Tiredness or weakness • Dizziness • Headache • Low blood pressure • Nausea <p>Neurologic problems</p> <p>Symptoms may include:</p> <ul style="list-style-type: none"> • Vomiting • Chills • Face swelling • Wheezing or trouble breathing • Skin rash <p>• Confusion or disorientation</p> <ul style="list-style-type: none"> • Loss of balance • Headache • Difficulty with facial movements, hearing, vision, or swallowing
<p>Medical information card</p> <p>Patient is receiving BLINCYTO® for the treatment of B-cell precursor acute lymphoblastic leukemia in first or second complete remission with minimal residual disease.</p> <p>BLINCYTO® (blinatumomab) for injection 35 mcg single-dose vial</p>	<p>IMPORTANT NOTE FOR HEALTH CARE PROFESSIONALS:</p> <p>Do not flush the BLINCYTO® infusion line or intravenous catheter, especially when changing infusion bags. Flushing when changing bags or at completion of infusion can result in excess dosage and complications.</p> <p>BLINCYTO® may cause serious side effects that can be severe, life-threatening, or fatal, including:</p>

Fold

Fold